



## Declaration by Dr Kyoko Higuchi

RECEIVED  
AUG 11 2003  
TECH CENTER 1600/2900

1. I am a graduate of the University of Tokyo from where I received my Ph.D. in 1985.
2. As a senior lecturer, I have taught and performed research in applied biology and chemistry in the Tokyo University of Agriculture.
3. I have worked on nicotianamine synthase (NAS) for many years. My publications in this field in the period 1994-2001 are listed in the attached annex.
4. I am named as an inventor for United States Patent Application No. 09/674,337. I have been asked to comment on the difficulties I encountered in purifying the NAS protein and cloning the NAS gene subsequent to performing the work described in Higuchi et al (1994) Plant and Soil 165, 173-9 (Higuchi et al 1994).
5. In Higuchi et al 1994, I reported partial purification of NAS and showed that NAS activity was induced under Fe-deficient condition in barley root. As mentioned in the final paragraph, left hand column, page 178 of Higuchi et al 1994. I then proceeded to attempt to clone NAS based on partial amino acid sequencing of the protein preparation described in Higuchi et al 1994. At this point I should mention that no NAS protein or related protein had previously been cloned. Therefore no sequence information was available to assist in the cloning of NAS.
6. I found that we could not obtain any N-terminal sequence information by subjecting the extract of Higuchi et al 1994 to amino acid sequencing. After many attempts to obtain sequence in this way failed, I attempted cloning of NAS based on an antibody based approach. Once we had obtained an antibody specific to NAS we could use this to purify NAS (for example using the antibody in an affinity column) or to screen

an expression library of clones from a plant that expresses NAS.

7. We generated a polyclonal antibody using the available extract that contained NAS. However, the antibody had only a weak reactivity to NAS protein, because the NAS extract we had used to generate the antibody was not pure enough. We found that the antibody was of no assistance in helping us to clone the NAS gene.
8. It is true that we subsequently did generate a high affinity antibody (as described in Higuchi et al (1999) Soil Sci. Plant Nutr., 45, 681-91. However this antibody was generated using a preparation of NAS obtained by following the procedure described in Higuchi. et al (1999) Plant Physiol., 119, 471-480, and not using the extract of Higuchi et al 1994.
9. I then attempted another approach to clone the NAS gene. We tried to purify NAS using based on its affinity for its substrate. A SAM analogue, S-adenosyl-homocysteine, was fixed to a column through which the NAS extract was passed. However, we found that elutions from the column either contained no protein or contained all of the protein that was present in the initial extract, and therefore affinity purification using the substrate proved unsuccessful.
10. I then attempted to purify NAS using an extract from another species. It had been reported that Basidiobolus Meristoporus, which is one of Zygomycete species, secretes nicotianamine. I tried to perform purification of NAS from Basidiobolus Meristoporus. However, the NAS activity of a crude extract obtained from Basidiobolus Meristoporus was very low. Further, I found that the medium in which Basidiobolus Meristoporus was cultured contained soybean protein and other unfavorable compounds which inhibited enzyme purification.
11. After this attempt, I decided that the best way forward would be to try to develop a

purification procedure which could be used to obtain NAS in a form in which it could be sequenced. As described above purification based on specific affinity purification was unsuccessful, and I was required to develop a procedure that purified NAS but not on the basis of sequence.

12. There are many reasons why protein sequencing may not be successful. Such reasons include low quantities of the protein to be sequenced and the presence of contaminating proteins and chemicals. In the present case a further complication was caused by the fact that the NAS protein is extremely sensitive to degradation. Degradation not only reduces the quantity of NAS but generates degradation products which also interfere with sequencing. In order to prevent degradation of NAS a protease inhibitor needs to be present in the NAS preparation during purification.
13. This means that purifying NAS is difficult because it is not simply a matter of subjecting it to additional purification steps until it is sufficiently pure. Too many purification steps result in loss and degradation of the NAS. Instead the development of a purification procedure requires the finding of a combination of purification steps which results in separation from contaminating proteins and chemicals but does not cause substantial loss or degradation of NAS.
14. After intensive effort on the part of me and my colleagues, and four years after the publication of Higuchi et al 1994, we finally developed a procedure which allowed us to obtain NAS in a form which could be sequenced. As can be seen in the attached table this procedure is much more complicated than the purification procedure used in Higuchi et al 1994, and was definitely not routine to develop. In particular, one reason why the new procedure may be superior to the purification procedure described in Higuchi et al 1994 is that the new procedure is better at removing impurities which have a similar molecular mass to NAS or which cause degradation

of NAS.

15. Upon cloning the gene we deduced that the original NAS extract of Higuchi et al 1994 had in fact consisted of seven isozymes. To some extent this may also have caused difficulties in NAS purification and sequencing. I believe the presence of multiple copies of NAS genes probably contributed to the difficulty in PCR amplification of NAS genes using a primer with sequence deduced from the N terminal amino acid sequence. In order to overcome these difficulties we resorted to using a primer based on EST sequence (as discussed in Example 5 of the present application).
16. Given the difficulties which I encountered in cloning the NAS gene I believe that this gene could not have been cloned by any routine means that was available at the time. As described above, we attempted all available techniques to clone the NAS gene, and none of them worked. It required the development of a special purification technique to obtain NAS protein in a form in which it could be sequenced.

Signed Kyoko Higuchi

Date 06/26/2003

### Annex

#### Publications by Kyoko Higuchi concerning nicotianamine synthase and its function in the period 1994-2001

Higuchi K., Kanazawa K., Nishizawa N K., Chino M., Mori S.

Purification and characterization of nicotianamine synthase from Fe deficient barley roots.

Plant and Soil, 1994, 165 (2) : 173-179.

Higuchi K., Nishizawa N K., Yamaguchi H., Roemheld V., Marschner H., Mori S.

Response of nicotianamine synthase activity to Fe deficiency in tobacco plants as compared with barley.

Journal of Experimental Botany, 1995, 46 (289) : 1061-1063.

Higuchi K., Nishizawa N., Roemheld V., Marschner H., Mori S.

Absence of nicotianamine synthase activity in the tomato mutant 'chloronerva'.

Journal of Plant Nutrition, 1996, 19 (8-9) : 1235-1239.

Higuchi K., Kanazawa K., Nishizawa N K., Mori S.

The role of nicotianamine synthase in response to Fe nutrition status in Gramineae.

Plant and Soil, 1996, 178 (2) : 171-177.

Higuchi K., Suzuki K., Nakanishi H., Yamaguchi H., Nishizawa N. K., Mori S.

Cloning of nicotianamine synthase genes, novel genes involved in the biosynthesis of phytosiderophores.

Plant Physiology, 1999, 119 (2) : 471-480.

Suzuki K., Higuchi K., Nakanishi H., Nishizawa N. K., Mori S.

Cloning of nicotianamine synthase genes from Arabidopsis thaliana.

Soil Science and Plant Nutrition, 1999, 45 (4) : 993-1002.

Higuchi K., Nakanishi H., Suzuki K., Nishizawa N. K. Mori S.

Presence of nicotianamine synthase isozymes and their homologues in the root of graminaceous plants.

Soil Science and Plant Nutrition, 1999, 45 (3) : 681-691.

Higuchi K., Watanabe S., Takahashi M., Kawasaki S., Nakanishi H., Nishizawa N. K., Mori S.

Nicotianamine synthase gene expression differs in barley and rice under Fe-deficient conditions.

Plant Journal, 2001, 25: 159-168.

Higuchi K., Tani M., Nakanishi H., Yoshiwara T., Goto F., Nishizawa N. K., Mori S.

The expression of a barley HvNAS1 nicotianamine synthase gene promoter-gus fusion gene in transgenic tobacco is induced by Fe-deficiency in roots.

Biosci. Biotechnol. Biochem., 2001, 65 (7) : 1692-1696.

Higuchi K., Takahashi M., Nakanishi H., Kawasaki S., Nishizawa N. K., Mori S.

Analysis of transgenic rice containing barley nicotianamine synthase gene.

Soil Science and Plant Nutrition, 2001, 47: 315-322.

## TABLE COMPARING PURIFICATION METHODS

Higuchi et al (1994)  
Plant and Soil  
165, 173-179

TSK gel Butyl-Toyopearl 650M  
(Fractogel TSK Butyl650M, Merck)  
column (1 x 11cm) equilibrated with  
starting buffer for hydrophobic  
chromatography synthase. NAS eluted  
with water containing 1% glycerol.

↓  
The enzyme fraction was equilibrated with  
the starting buffer (1 mM K-P, 0.1 M KCl,  
pH 8.0) for the hydroxylapatite chromato-  
graphy, concentrated, and applied to a hydroxylapatite 100~  
350 mesh (Nacalai Tesque Japan) column (1.5 x 2.7cm)  
equilibrated with the starting buffer. NAS was eluted with  
buffer A (4 mM K-P, 0.1 M KCl, pH 8.0 with KOH), and  
buffer B (8 mM K-P, 0.1 M KCl, pH 8.0).

↓  
SDS-PAGE was carried out at room temperature using  
12.5% acrylamide slab gels. After SDS PAGE, the gel was  
stained with 0.3 M CuCl<sub>2</sub> and was horizontally cut to several  
fractions nearby 30 Kda. Each fragment was destained  
with 0.25 M EDTA/0.25 M Tris(pH 9.0)

↓  
Polypeptides were eluted by electroelution  
with the buffer containing 7 mM Tris, 35 mM glycine and  
1% (v/v) glycerol.

(From Higuchi et al.,1994, Right hand column, page 174).

United States Patent Application No.  
09/674,337

TSK gel Butyl Toyopearl 650M column (10 ml bed volume  
per 100g of roots), equilibrated with the adsorption buffer  
and eluted with elution buffer [10 mM Tris, 1mM EDTA, 3  
mM DTT, 0.1 mM p-APMSF, 5% glyceroland 0.05% 3-  
[(3-chloramidopropyl) dimethylammonio] propanesulfonic  
acid (hereinafter called CHAPS), pH 8.0].

↓  
A hydroxapatite 100350 mesh (Nacalai Tesque),  
equilibrated with the adsorption buffer (1mM KP, 10 mM  
KCl, 3 mM DTT and 0.1 mM p-APMSF, pH 8.0), was  
prepared at 10 ml per protein 100 mg and the fractions  
containing nicotianamine synthase were loaded.  
Nicotianamine synthase was passed through without  
adsorption.

↓  
The passed through fraction was loaded onto TSK gel Butyl  
Toyopearl 650M column (1 ml bed volume per 10 mg of  
protein), and nicotianamine synthase was eluted as described  
above for the first step.

↓  
The active fraction was loaded onto a DEAE Sepharose FF  
column (5 ml bed volume per 25 mg of protein, Pharmacia)  
equilibrated with the adsorption buffer (20 mM Tris, 1 mM  
EDTA, 3 mM DTT, 0.1 mM p-APMSF and 0.05% CHAPS,  
pH 8.0) and eluted with stepwise gradient elution of  
potassium chloride concentration of 0.05 M, 0.1 M, 0.15 M  
and 0.2 M. Nicotianamine synthase was eluted at 0.15 M of  
KCl concentration.

↓  
The active fraction was loaded onto the Ether Toyopearl  
650M column (10 ml bed volume per 100g of roots),  
equilibrated with adsorption buffer [20 mM Tris, 1 mM  
EDTA, 3 mM DTT, 1.2 M (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> and 0.1 mM  
p-APMSF, pH 8.0]. Nicotianamine synthesis was not  
absorbed and passed through the column

↓  
The passed through fraction was loaded onto TSK gel Butyl  
Toyopearl 650M column and fractions containing  
nicotianamine synthase were eluted.

↓  
SDS-PAGE using 11% acrylamide slab gels. After  
SDS-PAGE the gel was stained with 0.3 M copper chloride  
and the separated bands were cut out. The gel fragments  
were destained with 0.25 M EDTA/0.25 M Tris (pH 9.0)  
and homogenized with the extraction buffer (1% SDS, 25  
mM Tris and 192 mM glycine). Each homogenate was  
electroeluted with SDS-free buffer (25 mM Tris and 192  
mM glycine) and peptide was recovered.

(From Examples 3 of United States Patent Application No.  
09/674,337)

	10	20	30	40	50	60
SEQ ID NO:1 HvNAS1 (barley)	MDAQ--	NKEVAALTEKTAGTQAAIA	LPSSLSPSPVDRLFT	DLVTACVPPSP	VDVTKLS	
SEQ ID NO:3 HvNAS2 (barley)	MAAQN--	QVDALVEKITGLHAAIAKL	PSLSPSPDVALFT	ELVTACVPPSP	VDVTKLG	
SEQ ID NO:5 HvNAS3 (barley)	MAAQN--	NNKDVAAALVEKITGLHAAIAKL	PSLSPSPDVALFT	ELVTACVPPSP	VDVTKLG	
SEQ ID NO:7 HvNAS4 (barley)	MDGQSE--	EVDALVQKITGLHAAIAKL	PSLSPSPDVALFT	DLVTACVPPSP	VDVTKLA	
SEQ ID NO:9 HvNAS5 (barley)	MEAEENG--	EVAALVEKITGLHAAIAKL	PSLSPSPDVALFT	ELVTACVPPSP	VDVTKLG	
SEQ ID NO:11 HvNAS6 (barley)	MDAQN--	EVDALVQKITGLHAAIAKL	PSLSPSPDVALFT	DLVTACVPPSP	VDVTKLG	
SEQ ID NO:13 HvNAS7 (barley)	MDAQSK--	EVDALVQKITGLHAAIAKL	PSLSPSPDVALFT	DLVTACVPPSP	VDVTKLA	
SEQ ID NO:15 OsNAS1 (rice)	MEAQNQ--	EVAALVEKITGLHAAIAKL	PSLSPSPDVALFT	DLVTACVPPSP	VDVTKLG	
SEQ ID NO:17 AtNAS1 (Arabido)	MACQNN----	LVVVKQIDLDYQDQISL	KSLKPSKNVDL	TFGQVSTCLP	TDNTDVTNMC	
SEQ ID NO:19 AtNAS2 (Arabido)	MACQNN----	LVVVKQIDLDYQDQISL	KSLKPSKNVDL	TFGQVSTCLP	TDNTDVTNMC	
SEQ ID NO:21 AtNAS3 (Arabido)	MGCQDE----	OLVQTHCDLYEKISKL	ESLKPSEDNV	LEKQVSTCLP	ENEDVTNMC	

	70	80	90	100	110	120
SEQ ID NO:1 HvNAS1 (barley)	-PEHQRMR	ALIRLCSA	AEGLKLAHYADLLAT	FDN---	PLDHLG	LPYYSNYNLSKLEY
SEQ ID NO:3 HvNAS2 (barley)	-PEAQEMR	ELIRLCSA	AEGLKLAHYSDMLA	AFDK---	PLDHLG	MPYYSNYNLSKLEY
SEQ ID NO:5 HvNAS3 (barley)	-PEAQEMR	ELIRLCSA	AEGLKLAHYSDMLA	AFDK---	PLDHLG	MPYYSNYNLSKLEY
SEQ ID NO:7 HvNAS4 (barley)	-PEAQAMR	ELIRLCSA	AEGLKLAHYSDMLA	AFDK---	PLDHLG	MPYYSNYNLSKLEY
SEQ ID NO:9 HvNAS5 (barley)	-PEAQEMR	DLIRLCSA	AEGLKLAHYSDMLA	AFDK---	PLDHLG	MPYYSNYNLSKLEY
SEQ ID NO:11 HvNAS6 (barley)	-PEAQEMR	ELIRLCSA	AEGLKLAHYSDMLA	AFDK---	PLDHLG	MPYYSNYNLSKLEY
SEQ ID NO:13 HvNAS7 (barley)	-PEAQAMR	ELIRLCSA	AEGLKLAHYSDMLA	AFDK---	PLDHLG	MPYYSNYNLSKLEY
SEQ ID NO:15 OsNAS1 (rice)	-PEAQAMR	ELIRLCSA	AEGLKLAHYSDMLA	AFDK---	PLDHLG	MPYYSNYNLSKLEY
SEQ ID NO:17 AtNAS1 (Arabido)	-EEVKDMR	ANLIRLCSA	AEGLKLAHYSDMLA	AFDK---	PLDHLG	MPYYSNYNLSKLEY
SEQ ID NO:19 AtNAS2 (Arabido)	DEKVKDMR	SHLIRLCSA	AEGLKLAHYSDMLA	AFDK---	PLDHLG	MPYYSNYNLSKLEY
SEQ ID NO:21 AtNAS3 (Arabido)	-DRVCEI	RLNLIKICG	LAEGHLENHPS	SLITSYQD---	NPLHHLN	IFYYNNYKLGKLEF

	130	140	150	160	170	180
SEQ ID NO:1 HvNAS1 (barley)	ELLARHVP-	GIAPARVAF	IGSGPLPFSS	VLAAHHP	ETQFDNYD	LCGAANERARKLFGA
SEQ ID NO:3 HvNAS2 (barley)	ELLARYVPG	GYRPARVAF	IGSGPLPFSS	FVLAARHLP	PTMFDNYD	LCGAANDRASKLFRA
SEQ ID NO:5 HvNAS3 (barley)	ELLARYVRR-	HRPARVAF	IGSGPLPFSS	FVLAARHLP	PTMFDNYD	LCGAANDRASKLFRA
SEQ ID NO:7 HvNAS4 (barley)	ELLARYVPG	RHRPARVAF	IGSGPLPFSS	FVLAARHLP	PTMFDNYD	LCGAANDRASKLFRA
SEQ ID NO:9 HvNAS5 (barley)	ELLARHVP-	GIAPARVAF	IGSGPLPFSS	VLAAHHP	ETQFDNYD	LCGAANDRASKLFRA
SEQ ID NO:11 HvNAS6 (barley)	ELLARYVPG	GIAPARVAF	IGSGPLPFSS	VLAAHHP	ETQFDNYD	LCGAANDRASKLFRA
SEQ ID NO:13 HvNAS7 (barley)	ELLARYVPG	GIAPARVAF	IGSGPLPFSS	VLAAHHP	ETQFDNYD	LCGAANDRASKLFRA
SEQ ID NO:15 OsNAS1 (rice)	DLIVRYVPG-	IASTRVAF	IGSGPLPFSS	VLAAHHP	ETQFDNYD	LCGAANDRASKLFRA
SEQ ID NO:17 AtNAS1 (Arabido)	DLISQHS-	--HVTETK	IAFVSGGPMPLT	SIVLAKF	HPENTFHN	EDIDSHANTLASNLVSR
SEQ ID NO:19 AtNAS2 (Arabido)	DLISQHTT-	HYETK	IAFVSGGPMPLT	SIVLAKF	HPENTFHN	EDIDSHANTLASNLVSR
SEQ ID NO:21 AtNAS3 (Arabido)	DLLEQNLG-	FVEKSVAF	IGSGPLPFSS	SIVLAKF	HPENTFHN	EDIDSHANTLASNLVSR

	190	200	210	220	230	240
SEQ ID NO:1 HvNAS1 (barley)	TADCVGARMS	FHTADVADLT	QELGAYDVVFLAALV	GMAAEK	KAKVIAHLGAHM	VEGASILV
SEQ ID NO:3 HvNAS2 (barley)	CRD-VGARMS	FHTADVADLT	QELGAYDVVFLAALV	GMAAEK	KAKVIAHLGAHM	VEGASILV
SEQ ID NO:5 HvNAS3 (barley)	CTD-VGARMS	FHTADVADLT	QELGAYDVVFLAALV	GMAAEK	KAKVIAHLGAHM	VEGASILV
SEQ ID NO:7 HvNAS4 (barley)	CKD-VGARMS	FHTADVADLT	QELGAYDVVFLAALV	GMAAEK	KAKVIAHLGAHM	VEGASILV
SEQ ID NO:9 HvNAS5 (barley)	ADEGVRSM	AFHTAEVDLT	QELGAYDVVFLAALV	GMAAEK	KAKVIAHLGAHM	VEGASILV
SEQ ID NO:11 HvNAS6 (barley)	CKD-VGARMS	FHTADVADLT	QELGAYDVVFLAALV	GMAAEK	KAKVIAHLGAHM	VEGASILV
SEQ ID NO:13 HvNAS7 (barley)	CKD-VGARMS	FHTADVADLT	QELGAYDVVFLAALV	GMAAEK	KAKVIAHLGAHM	VEGASILV
SEQ ID NO:15 OsNAS1 (rice)	ADEGLGARMA	FHTADVADLT	QELGAYDVVFLAALV	GMAAEK	KAKVIAHLGAHM	VEGASILV
SEQ ID NO:17 AtNAS1 (Arabido)	DPD-LSKRM	IFHTTOVLNATEAL	QDQDVVFLAALV	GMDKESKVK	KAIEHLEKHM	AFGAVLM
SEQ ID NO:19 AtNAS2 (Arabido)	CSD-LSKRM	IFHTTOVLNATEAL	QDQDVVFLAALV	GMDKESKVK	KAIEHLEKHM	AFGAVLM
SEQ ID NO:21 AtNAS3 (Arabido)	DPD-LSKRM	IFHTTOVLNATEAL	QDQDVVFLAALV	GMDKESKVK	KAIEHLEKHM	AFGAVLM

	250	260	270	280	290	300
SEQ ID NO:1 HvNAS1 (barley)	VRSAR-PRG	FLYPIVD	EDIRGGGF	EVLA	CHPE-GEVINS	SVIVARKA--VEAQ-L-SGP
SEQ ID NO:3 HvNAS2 (barley)	VRSAGARG	FLYPIVD	PDIGRGGGF	EVLA	CHPD-DDVNSV	IIAOKSKDVHADGLGSGR
SEQ ID NO:5 HvNAS3 (barley)	VRSAGARG	FLYPIVD	PDIGRGGGF	EVLA	CHPD-DDVNSV	IIAOKSKDVHADGLGSGR
SEQ ID NO:7 HvNAS4 (barley)	--ARHGARG	FLYPIVD	PDIGRGGGF	EVLA	CHPD-DDVNSV	IIAOKSKDVHADGLGSGR
SEQ ID NO:9 HvNAS5 (barley)	REALHGARG	FLYPIVD	PDIGRGGGF	EVLA	CHPD-DDVNSV	IIAOKSKDVHADGLGSGR
SEQ ID NO:11 HvNAS6 (barley)	V-RSAQARG	FLYPIVD	PDIGRGGGF	EVLA	CHPD-DDVNSV	IIAOKSKDVHADGLGSGR
SEQ ID NO:13 HvNAS7 (barley)	R-SAHGARG	FLYPIVD	PDIGRGGGF	EVLA	CHPD-DDVNSV	IIAOKSKDVHADGLGSGR
SEQ ID NO:15 OsNAS1 (rice)	VRTAHGARG	FLYPIVD	PDIGRGGGF	EVLA	CHPD-DDVNSV	IIAOKSKDVHADGLGSGR
SEQ ID NO:17 AtNAS1 (Arabido)	LRAHALR	FLYPIVD	SSDLK--GFQ	LTIIYHT	DDVNSV	IIAOKSKDVHADGLGSGR
SEQ ID NO:19 AtNAS2 (Arabido)	LRSAGHLR	FLYPIVD	SSDLK--GFQ	LTIIYHT	DDVNSV	IIAOKSKDVHADGLGSGR
SEQ ID NO:21 AtNAS3 (Arabido)	LRSAGHLR	FLYPIVD	SSDLK--GFQ	LTIIYHT	DDVNSV	IIAOKSKDVHADGLGSGR

	310	320	330	340
SEQ ID NO:1 HvNAS1 (barley)	QNGDAHARG-	AVPIVSPPCN	ESTKM-EASAL	-----EKSEELTAKELAF
SEQ ID NO:3 HvNAS2 (barley)	GAGGQYARG-	TVPVVSPPCN	RGEMVADVTQ	-----HKRDEEFANAEVAF
SEQ ID NO:5 HvNAS3 (barley)	GAGGQYARG-	TVPVVSPPCN	RGEMVADVTQ	-----HKRDEEFANAEVAF
SEQ ID NO:7 HvNAS4 (barley)	GRGGRYARG-	TVPVVSPPCN	RGEMVADVTQ	-----KREEFANAEVAF
SEQ ID NO:9 HvNAS5 (barley)	GRGGRYARG-	TVPVVSPPCN	RGEMVADVTQ	-----KREEFANAEVAF
SEQ ID NO:11 HvNAS6 (barley)	GRGGRYARG-	TVPVVSPPCN	RGEMVADVTQ	-----KREEFANAEVAF
SEQ ID NO:13 HvNAS7 (barley)	GCGGRYARG-	TVPVVSPPCN	RGEMVADVTQ	-----KREEFANAEVAF
SEQ ID NO:15 OsNAS1 (rice)	RDELADSR	GVVLPVVG	PPSTCKKVEASAVEK	-----AEBFAANKELSV
SEQ ID NO:17 AtNAS1 (Arabido)	-----TRG	-----CMF	PCNCSKIHAIMNNR	-----GKKNMIEEFSTIE
SEQ ID NO:19 AtNAS2 (Arabido)	-----IGR	-----CVV	MPNCSKVHAILNNR	-----GMEKNLIEEFSAIE
SEQ ID NO:21 AtNAS3 (Arabido)	-----PNS	-----CLL	PCNCSKTHAKMNKN	-----MMIEEFGARIEQLS